



DIAGNOSTIC AUTOMATION, INC.

21250 Califa Street, Suite 102 and 116, Woodland Hills, California 91367 USA

Tel: (818) 591-3030 Fax: (818) 591-8383

onestep@rapidtest.com

technicalsupport@rapidtest.com

www.rapidtest.com



See external label



2°C-8°C



96 tests



5140-8

Enzyme Immunoassay for the Quantitative Determination of Hazelnut in Food

Hazelnut

REF 5140-8

Sensitivity	0.33 PPM
Recovery (spiked samples)	83-101%
Incubation Time	60 min

GENERAL INFORMATION

Hazelnut (*Corylus avellana*) belongs to the birch plants. With 13 % the fraction of proteins in hazel-nuts is high. Many of these proteins are known for being allergenic, such as Cor a 9 and Cor a 11. Most of them are very heat resistant making them stable to different production processes. For this reason hazelnut represents an important food allergen. For hazelnut allergic persons hidden hazelnut allergens in food are a critical problem. Already very low amounts of hazelnuts can cause allergic reactions, which may lead to anaphylactic shock in severe cases. Because of this, hazelnut allergic persons must strictly avoid the consumption of hazelnuts or hazelnut containing food. Cross contamination, mostly in consequence of the production process is often noticed. The chocolate production process is a representative example. This explains why in many cases the existence of hazelnut residues in foods cannot be excluded. For this reason sensitive detection systems for hazelnut residues in foodstuffs are required.

The **DAI ELISA** represents a highly sensitive detection system and is particularly capable of the quantification of hazelnut residues in cookies, cereals, ice cream and chocolate.

PRINCIPLE OF THE TEST

The DAI Hazelnut quantitative test is based on the principle of the enzyme linked immunosorbent assay. An antibody directed against hazelnut proteins is bound on the surface of a microtiter plate. Hazelnut containing samples or standards are given in-to the wells of the microtiter plate. After 20 minutes incubation at room temperature, the wells are washed with diluted washing solution to remove unbound

Diagnostic Automation/ Cortez Diagnostics, Inc.
 21250 Califa St, Suite 102 and 116, Woodland Hills, CA 91367 USA Phone: 818-591-3030, Fax: 818-591-8383
 Email: onestep@rapidtest.com Website: www.rapidtest.com

material. A peroxidase conjugated second antibody directed against hazelnut proteins is given into the wells and after 20 minutes of incubation the plate is washed again. A substrate solution is added and incubated for 20 minutes, resulting in the development of a blue color. The color development is inhibited by the addition of a stop solution, and the color turns yellow. The yellow color is measured photometrically at 450 nm. The concentration of hazelnut is directly proportional to the color intensity of the test sample.

PRECAUTIONS

Full compliance of the following good laboratory practices (GLP) will determine the reliability of the results:

1. Prior to beginning the assay procedure, bring all reagents to room temperature (20-25°C).
2. All reagents should be mixed by gentle inversion or swirling prior to use. Do not induce foaming.
3. Once the assay has been started, all subsequent steps should be completed without interruption and within the recommended time limits.
4. Replace caps in all the reagents immediately after use. Do not interchange vial stoppers.
5. Use a separate disposable tip for each specimen to prevent cross-contamination.
6. All specimens and standards should be run at the same time, so that all conditions of testing are the same.
7. Do not mix components from different batches.
8. Do not use reagents after expiration date.
9. Check both precision and accuracy of the laboratory equipment used during the procedure (micropipets, ELISA reader etc.).

HEALTH AND SAFETY INSTRUCTIONS

1. Do not smoke or eat or drink or pipet by mouth in the laboratory.
2. Wear disposable gloves whenever handling patient specimens.
3. Avoid contact of substrate and stop solution with skin and mucosa (possible irritation, burn or toxicity hazard). In case of contact, rinse the affected zone with plenty of water.
4. Handling and disposal of chemical products must be done according to good laboratory practices (GLP).

REAGENTS

The kit contains reagents for 96 determinations. They have to be stored at 2-8°C. Expiry data are found on the labels of the bottles and the outer package.

1. Microtiter plate consisting of 12 strips with 8 breakable wells each, coated with anti-hazelnut antibodies.
2. Hazelnut Standards (0; 1; 4; 10; 40 ppm of hazelnut): 5 vials with 1.0 mL each, dyed red, ready-to-use
3. Conjugate (anti-hazelnut-peroxidase): 15 mL, dyed red, ready-to-use.
4. Substrate Solution (TMB): 15 mL, ready-to-use.
5. Stop Solution (0.5 M H₂SO₄): 15 mL, ready-to-use.
6. Extraction and sample dilution buffer (Tris): 2 x 120 mL as 10x concentrate, dyed red. Dilute 1+9 with distilled water. Stored at 4°C the diluted buffer is stable for at least one week. If during the cold storage crystals precipitate, the concentrate should be warmed up to 37°C for 15 minutes.
7. Washing Solution (PBS + Tween 20): 60 mL as 10x concentrate. Dilute 1+9 with distilled water. Stored at 4°C the diluted buffer is stable for at least 4 weeks. If during the cold storage crystals precipitate, the concentrate should be warmed up to 37°C for 15 minutes.
8. Plastic bag to store unused microtiter strips.
9. Instruction Manual.

Additional Instrumentation and Reagents (not provided)

Instrumentation

- 100 - 1000 μ L micropipets
- Volumetric flask
- Analytical balance
- Mortar, mixer
- Water bath
- Centrifuge
- ELISA reader (450 nm)

Reagents

- Double distilled water

SAMPLE PREPARATION

Due to high risk of cross-contamination all applied instruments like applicator, mortar, glass vials etc. have to be cleaned thoroughly before and after each sample. Hazelnut proteins adhere very strongly to different surfaces. In certain cases they can resist a common dishwasher cleaning. To identify possible cross-contamination caused by previous extractions it is strongly recommended to note the sequence of the extractions.

The following sample preparation should be applied for all kinds of samples:

1. To maximize homogeneity and representativeness of the sample drawing, a minimum of 5 g sample should be pulverized finely in a mortar, impact mill etc.
2. 1 g of the homogenized mixture is suspended in 20 mL of pre-diluted extraction buffer. Afterwards the suspension is incubated for 15 min in a preheated water bath at 60°C. To ensure good homogeneity, the samples should be shaken every two minutes.
3. The samples are centrifuged for 10 minutes at 2000 g. If it is not possible to separate the supernatant from the precipitate completely, the suspension should be filtrated if necessary.
4. 100 μ L of particle-free solution are applied per well. If the results of a sample are out of the measuring range, further dilution with the pre-diluted extraction and sample dilution buffer is necessary. The additional dilution has to be considered when calculating the concentration.

PROCEDURE

The washing solution is supplied as 10x concentrate and has to be **diluted** 1+9 with double distilled water before use.

In any case the **ready-to-use** standards provided should be determined twofold. When samples in great quantities are determined, the standards should be pipetted once before the samples and once after the samples. For final interpretation the arithmetic mean is used for calculation.

In consideration of GLP and quality control requirements a duplicate measurement of samples is recommended.

The procedure is according to the following scheme:

1. Prepare samples as described above.

Diagnostic Automation/ Cortez Diagnostics, Inc.
21250 Califa St, Suite 102 and 116, Woodland Hills, CA 91367 USA Phone: 818-591-3030, Fax: 818-591-8383
Email: onestep@rapidtest.com Website: www.rapidtest.com

2. Pipet 100 μ L ready-to-use standards or prepared samples in duplicate into the appropriate wells of the microtiter plate.
3. Incubate for 20 minutes at room temperature.
4. Wash the plate three times as follows: Discard the contents of the wells (dump or aspirate). Pipet 300 μ L of diluted washing solution into each well. After the third repetition empty the wells again and remove residual liquid by striking the plate against a paper towel. The wash procedure is critical. Insufficient washing will result in poor precision and falsely elevated absorbencies.
5. Pipet 100 μ L of conjugate (anti-hazelnut-peroxidase) into each well.
6. Incubate for 20 minutes at room temperature.
7. Wash the plate as outlined in 4.
8. Pipet 100 μ L of substrate solution into each well.
9. Allow the reaction to develop in the dark (e.g. cupboard or drawer; the chromogen is light-sensitive) for 20 minutes at room temperature.
10. Stop enzyme reaction by adding 100 μ L of stop solution (0.5 M H₂SO₄) into each well. The blue color will turn yellow upon addition.
11. After thorough mixing, measure absorbance at 450 nm (reference wavelength 620 nm), using an ELISA reader. The color is stable for 30 minutes.

CALCULATION OF RESULTS

The ready-to-use standards are prepared for a direct determination of sample concentrations. The dilution of samples in the extraction process as described in the above stated sample preparation procedure is already considered. Additional dilution due to high sample concentration has to be accounted for.

1. Calculate the average optical density (OD 450 nm) for each set of reference standards or samples.
2. Construct a standard curve by plotting the mean optical density obtained for each reference standard against its concentration in ppm on semi-log graph paper with the optical density on the vertical (y) axis and the concentration on the horizontal (x) axis. Alternatively the evaluation can be carried out by software. In this case the 4-parameter method should be preferred.
3. Using the mean optical density value for each sample, determine the corresponding concentration of hazelnut in ppm from the standard curve. Depending on experience and/or the availability of computer capability, other methods of data reduction may be employed.

TYPICAL STANDARD VALUES

The following table contains an example for a typical standard curve. The binding is calculated as percent of the absorption of the 40 ppm standard. These values are only an example and should not be used instead of the standard curve which has to be measured in every new test.

Hazelnut (ppm)	% binding of 40 ppm
40	100
10	41
4	23
1	9
0	5

PERFORMANCE

Sensitivity

The limit of detection (LOD) of the DAI Hazelnut test is 0.3 ppm.

The limit of quantification (LOQ) of the DAI Hazelnut test is 1 ppm.

Due to the variety of sample matrices and their influence on the blank, results less than the LOQ should be treated as negative.

Cross-reactivity

For the following foods no cross-reactivity could be detected:

Wheat	Soy	Brazil nut
Barley	Poppy seed	Pistachio
Rye	Sunflower seed	Macadamia nut
Oats	Pumpkin seed	Chestnut
Buckwheat	Pine nuts	Cocoa
Corn	Cashew nut	Dried milk
Rice	Sesame	Gluten
Pea	Peanut	Lecithin
Chickpea	Almond	Gelatin
Bean	Coconut	Apple

The following cross reactions were determined:

Walnut	< 0.0022%
--------	-----------

Precision

Intra-assay Precision	7-12%
Inter-assay Precision	3-13%

Linearity

The serial dilution of spiked samples (cookies, cereals, ice cream and chocolate) resulted in a dilution linearity of 87% - 121%.

Recovery

Mean recovery was determined by spiking samples with different amounts of hazelnut:

Cookies	101%
Cereals	99%
Ice cream	90%
Chocolate	83%

REFERENCES

1. Wigotzki M, et al. (2000) – Influence of varieties, storage and heat treatment on IgE-binding proteins in hazelnuts (*Corylus avellana*). Food and Agric Immunol, 12:217-29
2. Le TM, et al. (2008) – Reported food allergy to peanut, tree nuts and fruit: comparison of clinical manifestations, prescription of medication and impact on daily life. Allergy, 63(7):910-6
3. Luettkopf D, et al. (2001) – Comparison of four variants of major allergen in hazelnut (*Corylus avellana*) Cor a 1.04 with the major hazel pollen allergen Cor a 1.01. Mol Immunol, 38:515-25
4. Ballmer-Weber BK, et al. (2008) – Predictive value of the sulfidoleukotriene release assay in oral allergy syndrome to celery, hazelnut, and carrot. J Investig Allergol Clin Immunol, 18(2):93-99
5. Ben Rejeb S, et al. (2005) – Multi-allergen screening immunoassay for the detection of protein markers of peanut and four tree nuts in chocolate. Food Addit Contam, 22(8):709-15
6. Lauer I, et al. (2004) – Hazelnut (*Corylus avellana*) vicilin Cor a 11: molecular characterization of a glycoprotein and its allergenic activity. Biochem J, 383:327-334
7. Holzhauser T, et al. (2002) – Detection of potentially allergenic hazelnut (*Corylus avellana*) residues in food: a comparative study with DNA PCR-ELISA and protein sandwich-ELISA. J Agric Food Chem, 50(21):5808-15
8. Kiening M, et al. (2005) – Sandwich immunoassays for the determination of peanut and hazelnut traces in foods. J Agric Food Chem, 53(9):3321-7
9. Holzhauser T, et al. (1999) – Quantitative sandwich ELISA for determination of traces of hazelnut (*Corylus avellana*) protein in complex food matrixes. J Agric Food Chem, 47(10):4209-18
10. Akkerdaas JH, et al. (2004) – A novel approach for the detection of potentially hazardous pepsin stable hazelnut proteins as contaminants in chocolate-based food. J Agric Food Chem, 52(25):7726-31

Date Adopted	2017-3-21
REF 5140-8	DA-Hazelnut



DIAGNOSTIC AUTOMATION, INC.

21250 Califa Street, Suite 102 and 116, Woodland Hills, California 91367 USA

Tel: (818) 591-3030 Fax: (818) 591-8383

ISO 13485-2003



Revision Date: 2015-Mar-4